

# Multi-phase transport model for heavy ion collisions at RHIC

Z. Lin<sup>a</sup> S. Pal<sup>a</sup> C. M. Ko<sup>a</sup> B. A. Li<sup>b</sup> B. Zhang<sup>b</sup>

<sup>a</sup>*Cyclotron Institute and Physics Department, Texas A&M University, College Station, Texas 77843*

<sup>b</sup>*Department of Chemistry and Physics, Arkansas State University, State University, Arkansas 72467*

---

*Presented by: Z. Lin*

---

## Abstract

To study heavy ion collisions at RHIC, we have developed a multi-phase transport (AMPT) model [1] by replacing jet quenching in the HIJING model with explicit parton cascade based on the ZPC model and including final-state hadronic interactions using the ART model. We find that a larger parton-parton cross section than used in the default ZPC model is required to reproduce the total charged particle multiplicity near mid-rapidity measured by the PHOBOS collaboration at RHIC for Au+Au collisions at center-of-mass energies of 56 and 130 AGeV. We also find that final-state hadronic interactions reduce the anti-proton to proton ratio due to annihilation. The AMPT model further allows us to study the relative importance of the partonic and hadronic stages on strangeness production,  $J/\psi$  suppression [2] and elliptic flow [3].

[1] B. Zhang, C.M. Ko, B.A. Li, and Z. Lin, Phys. Rev. C61, 067901 (2000). [2] B. Zhang, C.M. Ko, B.A. Li, Z. Lin, and B.H. Sa, Phys. Rev. C62, 054905 (2000). [3] B. Zhang, C.M. Ko, and M. Gyulassy, Phys. Lett. B455, 45 (1999).

---